

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-12532: Mouse Monoclonal Antibody to CK17(Clone :BS55)

Clonality: Monoclonal

Clone Name: BS55
Application: IHC
Reactivity: Human
Gene: KRT17
Gene ID: 3872
Uniprot ID: Q04695

Alternative Name: CK17, PC, K17, PC2, PCHC1, KRT17

**Isotype:** Mouse IgG2b

## **Description**

CK17, also known as KRT17, it is the type I intermediate filament chain keratin 17. It is found in nail beds, hair follicles, sebaceous glands, and other epidermal appendages. Mutations in this gene lead to Jackson-Lawler type pachyonychia congenita and steatocystoma multiplex. May play a role in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial "stem cells". May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation. Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state. Modulates the function of TNF-alpha in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway. Involved in tissue repair.

## **Product Info**

**Amount:** 0.1 ml / 0.5 ml

**Content:** TRIS with 0.03% sodium azide, pH7.2

**Storage condition :** Store at 4°C

## **Application Note**

Immunohistochemical Analysis:-1:200



Figure-1: Tonsil section has been stained using CK17 antibody (Clone: BS55) with 1:200 dilution. Epithelia of tonsil are stained intensively.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Figure-2: Tonsil section has been stained using CK17 antibody (Clone: BS55) with 1:200 dilution. Epithelia of tonsil are stained intensively.



Figure-3: Skin section has been stained using CK17 antibody (Clone: BS55) with 1:200 dilution. Epidermis of the skin stained with permanent red chromogen.