

# 10-3003-NALE: NALE™ Monoclonal Antibody to TLR9 (Clone: ABM1C51)

Clonality :	Monoclonal
Clone Name :	ABM1C51
Application :	IHC,FACS,WB
Reactivity :	Mouse,Human
Gene :	TLR9
Gene ID :	54106
Uniprot ID :	Q9NR96
Format :	Azide Free,Purified
Alternative Name :	TLR9,UNQ5798/PRO19605
Isotype :	Mouse IgG1 Kappa
Immunogen Information	A partial length recombinant TLR9 protein (amino acids 100-290) was used as the immunogen for the antibody.

## Description

TLR9, a member of toll-like receptor family are central to the innate immunity by identifying pathogen associated molecular patterns (PAMPs). TLR9 identify unmethylated CpG dinucleotides present in bacterial DNA leading to NF-kB activation.

#### **Product Info**

Amount :	100 µg
Purification :	Protein G Chromatography
Content :	25 μg in 50 μl/100 μg in 200 μl PBS containing No Azide, low endotoxin (0.1EU/1ug).
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### **Application Note**

Western blot analysis: 2-4 µg/ml, FACS analysis: 0.5 µg/10^6 cells, Immunohistochemical analysis: 5 µg/ml




Fig-1: Immunohistochemical analysis of TLR9 in human stomach tissue using TLR9 antibody (Clone: ABM1C51) at 5  $\mu$ g/ml.

# **w** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Fig-2: Western blot analysis of TLR9. Anti- TLR9 antibody (Clone: ABM1C51) was used at 2  $\mu$ g/ml on (1) Raji and (2) EL4 lysates.

Fig-3: Intracellular flow analysis of TLR9 in human PBMC (Lymphocytes) using 0.5  $\mu$ g/10^6 cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary.

Fig-4: Intracellular flow analysis of TLR9 in Raji cells using  $0.5 \ \mu g/10^6$  cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary antibody.

Fig-5 : Immunohistochemical analysis of TLR9 in human Liver tissue using TLR9 antibody (Clone: ABM1C51) at 20  $\mu$ g/ml.