

39-1018: Anti-CDC6 Monoclonal Antibody (Clone: DCS-180)

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | DCS-180 |
| Application : | WB |
| Reactivity : | Human |
| Gene : | Cdc6 |
| Gene ID : | 23834 |
| Uniprot ID : | O89033 |
| Alternative Name : | Cell division control protein 6 homolog; CDC6-related protein; p62(cdc6); Cdc6 |
| Isotype : | Mouse IgG1 |
| Immunogen Information : | Recombinant human Cdc6. |

Description

In yeasts, Cdc6(*Saccharomyces cerevisiae*) and Cdc18(*Schizosaccharomyces pombe*) associate with the origin recognition complex(ORC) proteins to render cells competent for DNA replication. Cdc6 is overexpressed in human cancers, where it has a critical regulatory role in addition to DNA replication. Yan et al.(1998) showed that Cdc6 is expressed selectively in proliferating but not quiescent mammalian cells, both in culture and within tissues in intact animals. During the transition from a growth-arrested to a proliferative state, transcription of mammalian Cdc6 is regulated by E2F proteins, as revealed by a functional analysis of the human Cdc6 promoter and by the ability of exogenously expressed E2F proteins to stimulate the endogenous Cdc6 gene. They conclude that expression of human Cdc6 is regulated in response to mitogenic signals through transcriptional control mechanisms involving E2F proteins, and that Cdc6 is required for initiation of DNA replication in mammalian cells.

Product Info

| | |
|----------------------------|--|
| Amount : | 100 µg/vial |
| Purification : | Ascites |
| Content : | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN ₃ as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100ug/ml. |
| Storage condition : | At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing. |

Application Note

Western blot : 1-2µg/ml