

## 39-1037: Anti-GAP43 Monoclonal Antibody (Clone: GAP-7B10)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	GAP-7B10
<b>Application :</b>	WB,IHC-P,IHC-F
<b>Reactivity :</b>	Human
<b>Gene :</b>	Gap43
<b>Gene ID :</b>	29423
<b>Uniprot ID :</b>	P07936
<b>Alternative Name :</b>	Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Protein F1; Gap43
<b>Isotype :</b>	Mouse IgG2a
<b>Immunogen Information :</b>	GAP-43 from neonatal rat forebrain membranes.

### Description

GAP43 is expressed by developing and regenerating neurons, and to a lesser extent, reactive glial cells. It is used widely to specifically label injured neurons and to score neuronal regeneration. GAP43 is also a neuronal growth cone protein thought to be involved in pathfinding. GAP43 is considered to be a crucial component of an effective regenerative response in the nervous system.

### Product Info

<b>Amount :</b>	100 µg/vial
<b>Purification :</b>	Ascites
<b>Content :</b>	Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN <sub>3</sub> as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100ug/ml.
<b>Storage condition :</b>	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

### Application Note

Western blot : 0.5-1 µg/ml; Immunohistochemistry(Paraffin-embedded Section) : 1-2 µg/ml;  
Immunohistochemistry(Frozen Section) : 1-2 µg/ml




Figure 1: Anti-GAP43 monoclonal antibody(39-1037). IHC(P): Rat Cardiac Muscle Tissue.