

39-2053: Anti-KCA3.1 Polyclonal Antibody

Clonality :	Polyclonal
Application :	WB
Reactivity :	Human
Gene :	KCNN4
Gene ID :	3783
Uniprot ID :	O15554
Alternative Name :	Intermediate conductance calcium-activated potassium channel protein 4; SK4; SKCa 4; SKCa4; IKCa1; IK1; KCa3.1; KCa4; Putative Gardos channel; KCNN4; IK1, IKCA1, KCA4, SK4
Isotype :	Rabbit IgG

Description

Intermediate conductance calcium-activated potassium channel protein 1(KCNN4, Kca3.1) is part of a potentially heterotetrameric voltage-independent potassium channel that is activated by intracellular calcium. Activation is followed by membrane hyperpolarization, which promotes calcium influx. KCNN4 may be part of the predominant calcium-activated potassium channel in T-lymphocytes. This gene is similar to other KCNN family potassium channel genes, but it differs enough to possibly be considered as part of a new subfamily.

Product Info

Amount :	100 µg/vial
Purification :	Immunogen affinity purified.
Content :	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ . Reconstitute : Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Storage condition :	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Application Note

Western blot : 0.1-0.5µg/ml; Immunohistochemistry(Paraffin-embedded Section) : 0.5-1µg/ml




Figure 1: Anti-KCNN antibody(39-2053). Western blotting: Lanes: Anti KCNN 39-2053 at 0.5ug/ml. WB: HUT Whole Cell Lysate at 40ug. Predicted band size: 60 kDa. Observed band size: 60 kDa.