

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 39-2054: Anti-NMDAR2A Polyclonal Antibody

Clonality: Polyclonal

Application: WB Reactivity: Human Gene: GRIN2A Gene ID: 2903 **Uniprot ID:** 012879

Glutamate receptor ionotropic, NMDA 2A; GluN2A; Glutamate [NMDA] receptor subunit epsilon-1; N-**Alternative Name:** 

methyl D-aspartate receptor subtype 2A; NMDAR2A; NR2A; hNR2A; GRIN2A; NMDAR2A

Isotype: Rabbit IgG

## **Description**

GRIN2A is also known as N-methyl-D-aspartate receptor channel, subunit epsilon-1(NMDAR2A). This gene encodes a member of the glutamate-gated ion channel protein family. The encoded protein is an N-methyl-D-aspartate (NMDA) receptor subunit. NMDA receptors are both ligand-gated and voltage-dependent, and are involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. These receptors are permeable to calcium ions, and activation results in a calcium influx into post-synaptic cells, which results in the activation of several signaling cascades. Disruption of this gene is associated with focal epilepsy and speech disorder with or without mental retardation. Alternative splicing results in multiple transcript variants.

## **Product Info**

Amount: 100 μg/vial

**Purification:** Immunogen affinity purified.

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3. Content:

Reconstitute: Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and Storage condition:

stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

## **Application Note**

Western blot: 0.1-0.5µg/ml; Immunohistochemistry(Paraffin-embedded Section): 0.5-1µg/ml




Figure 1: Anti-NMDAR2A antibody(39-2054). Western blotting: Lanes: Anti NMDAR2A(39-2054) at 0.5ug/ml, Lane 1: Rat Brain Tissue Lysate at 50ug, Lane 2: Mouse Brain Tissue Lysate at 50ug. Predicted band size: 165 kDa. Observed band size: 165 kDa.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Anti-NMDAR2A antibody(39-2054). IHC(P): Rat Brain Tissue.