

39-2092: Anti-FGF8 Polyclonal Antibody

Clonality :	Polyclonal
Application :	WB
Reactivity :	Human
Gene :	FGF8
Gene ID :	2253
Uniprot ID :	P55075
Alternative Name :	Fibroblast growth factor 8; FGF-8; Androgen-induced growth factor; AIGF; Heparin-binding growth factor 8; HBGF-8; FGF8; AIGF
Isotype :	Rabbit IgG

Description

Fibroblast growth factor 8 (androgen-induced), also known as FGF8 or AIGF, is a human gene which maps to 10q24. The protein encoded by this gene are secreted proteins that interact with FGF tyrosine kinase receptors to mediate growth and development. This protein is known to be a factor that supports androgen and anchorage independent growth of mammary tumor cells. Overexpression of this gene has been shown to increase tumor growth and angiogenesis. The temporal and spatial patterns of this gene expression suggest that FGF8 is involved in gastrulation, regionalization of the brain, and organogenesis of the limb and face as an embryonic epithelial factor. The adult expression of FGF8 is restricted to gonads, including testes and ovaries. FGF8 stimulated growth of human prostate carcinoma cells and mouse fibroblasts and mammary carcinoma cells in a dose-dependent manner.

Product Info

Amount :	100 µg/vial
Purification :	Immunogen affinity purified.
Content :	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ . Reconstitute : Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Storage condition :	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Application Note

Western blot : 0.1-0.5µg/ml; Immunohistochemistry(Paraffin-embedded Section) : 0.5-1µg/ml




Figure 1: Anti-FGF8 antibody(39-2092). Western blotting: Rat Ovary Tissue Lysate.




Figure 2: Anti-FGF8 antibody(39-2092). IHC(P): Rat Ovary Tissue.