

## 39-2128: Anti-Integrin beta 4 Polyclonal Antibody(Discontinued)

|                           |                                      |
|---------------------------|--------------------------------------|
| <b>Clonality :</b>        | Polyclonal                           |
| <b>Application :</b>      | WB                                   |
| <b>Reactivity :</b>       | Human                                |
| <b>Gene :</b>             | ITGB4                                |
| <b>Gene ID :</b>          | 3691                                 |
| <b>Uniprot ID :</b>       | P16144                               |
| <b>Alternative Name :</b> | Integrin beta-4; GP150; CD104; ITGB4 |
| <b>Isotype :</b>          | Rabbit IgG                           |

### Description

ITGB4(Integrin, beta-4), also known as CD104(Cluster of Differentiation 104), is a human gene. The gene encodes the integrin beta 4 subunits, a receptor for the laminins. This subunit tends to associate with alpha 6 subunits and is likely to play a pivotal role in the biology of invasive carcinoma. The ITGB4 gene is mapped on 17q25.1. Using expression profiling, Yang et al. found that ITGB4 was upregulated 6-fold by ZKSCAN3 in transfected human colon cancer cells compared with parental cells. They confirmed that ZKSCAN3 bound the promoter of ITGB4 in vitro and in vivo. ITGB4 knockdown by short hairpin RNA countered ZKSCAN3-augmented anchorage-independent colony formation in the colon cancer cell lines. The integrin beta-4 subunit is characterized by an unusually long cytoplasmic domain that harbors 4 fibronectin type III(FNIII) repeats, residing in 2 pairs separated by a connecting segment.

### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 100 µg/vial  |
| <b>Purification :</b>      | Immunogen affinity purified.   |
| <b>Content :</b>           | Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg NaN <sub>3</sub> .<br>Reconstitute : Add 0.2ml of distilled water will yield a concentration of 500ug/ml. |
| <b>Storage condition :</b> | At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.                                      |

### Application Note

Western blot : 0.1-0.5µg/ml; Immunohistochemistry(Paraffin-embedded Section) : 0.5-1µg/ml




Figure 1: Anti-ITGB4 antibody(39-2128). Western blotting: Lanes: Anti ITGB4(39-2128) at 0.5ug/ml. WB: A431 Whole Cell Lysate at 40ug. Predicted band size: 200 kDa. Observed band size: 200 kDa.