

39-2159: Anti-IL18 Polyclonal Antibody

Clonality :	Polyclonal
Application :	WB
Reactivity :	Rat
Gene :	Il18
Gene ID :	29197
Uniprot ID :	P97636
Alternative Name :	Interleukin-18; IL-18; Interferon gamma-inducing factor; IFN-gamma-inducing factor; Interleukin-1 gamma; IL-1 gamma; Il18; Igif
Isotype :	Rabbit IgG
Immunogen Information :	E.coli-derived rat IL18 recombinant protein (Position: H37-S194). Rat IL18 shares 64% and 91% amino acid (aa) sequence identity with human and mouse IL18, respectively.

Description

Interleukin (IL)-18, also called Interferon-gamma-inducing factor (IGIF), augments natural killer (NK) activity in spleen cells. The gene encodes a precursor protein of 192 amino acids and a mature protein of 157 amino acids. IL18 is a recently discovered cytokine that modulates both T helper type 1 (Th1) and Th2 responses. IL18 is a potent proinflammatory cytokine with potential atherogenic properties. It is highly expressed in the atherosclerotic plaques compared with control normal arteries and is localized mainly in plaque macrophages.

Product Info

Amount :	100 µg/vial
Purification :	Immunogen affinity purified.
Content :	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . Reconstitute : Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Storage condition :	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Application Note

Western blot : 0.1-0.5 µg/ml




Figure 1: Anti-IL-18 antibody(39-2159). Lanes: Anti-IL-18(39-2159) at 0.5ug/ml. Lane 1: Rat Spleen Tissue Lysate at 40ug, Lane 2: Rat Thymus Tissue Lysate at 40ug, Lane 3: PC12 Whole Cell Lysate at 40ug. Predicted band size: 22 kDa. Observed band size: 22 kDa.