

43-1007: Monoclonal Antibody to Mouse CD80 (B7-1) NALE™ Purified (Clone: 16-10A1)

Clonality :	Monoclonal
Clone Name :	16-10A1
Application :	FACS
Reactivity :	Mouse
Gene :	Cd80
Gene ID :	12519
Uniprot ID :	Q00609
Alternative Name :	Activation B7-1 antigen, B7
Isotype :	Armenian Hamster IgG

Description

The 16-10A1 antibody reacts with mouse CD80, also known as B7-1, a 55 kDa type I transmembrane protein ligand for CD152 (CTLA-4) and for CD28, a co-stimulatory receptor for the T cell receptor (TCR). CD28 also binds a second B7 ligand known as CD86 (B7-2). Both CD80 and CD86 are expressed on activated B cells and antigen-presenting cells. These ligands trigger CD28 signaling in concert with TCR activation to drive T cell proliferation, induce high-level expression of IL-2, impart resistance to apoptosis, and enhance T cell cytotoxicity. The interaction / co-stimulatory signaling between the B7 ligands and CD28 or CTLA-4 provides crucial communication between T cells and B cells or APCs to coordinate the adaptive immune response.

Product Info

Amount :	100 µg
Content :	2.0 mg/ml in Phosphate-buffered aqueous solution, pH7.2.
Storage condition :	The product should be stored undiluted at 4°C. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography. The endotoxin level is determined by LAL test to be less than 0.01 EU/µg of the protein.

Application Note

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. It is recommended that the reagent be titrated for optimal performance for each application.