

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-3549: Monoclonal Antibody to human Macrophages(Discontinued)

Clone Name: 25F9
Application: FACS
Reactivity: Human
Isotype: Mouse IgG1

## **Description**

The monoclonal antibody 25F9 recognises a protein of 86 kD on the cell surface and within the cytoplasm of mature macrophages. The antibody is associated with fully differentiated tissue macrophages both in normal and in diseased tissues, independently of the presence or absence of inflammation. The antigen is absent on freshly isolated monocytes and other blood cells. After 6 to 7 days culture human monocytes become positive. Some melanoma and carcinoma cell lines are also positive. Furthermore the monoclonal antibody 25F9 cross reacts with a subpopulation of macrophages of rhesus monkey, pig alveolar macrophages and Kupffer cells. The monoclonal antibody 25F9 is very useful for macrophage phenotyping.

## **Product Info**

Amount: Monoclonal Antibody to human Macrophages(Discontinued) / 500 µg

Content: 0.5 mg 0.2 µm filtered antibody solution in PBS, containing 0.1% bovine serum albumin and

0.02% sodium azide.

**Storage condition :** Product should be stored at 4°C. Under recommended storage conditions, product is stable for

one year.

## **Application Note**

For immunohistology, flow cytometry and Western blotting, dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.