

32-7008: Recombinant Human Interleukin-6/IL-6

Gene : IL6
Gene ID : 3569
Uniprot ID : P05231

Description

Source: E.coli.
MW :20.9kD.

Recombinant Human Interleukin-6 is produced by our E.coli expression system and the target gene encoding Pro29-Met212 is expressed. Cytokines of the IL6/GCSF/MGF family are glycoproteins of about 170 to 180 amino acid residues that contain four conserved cysteine residues involved in two disulfide bonds. They have a compact, globular fold (similar to other interleukins), stabilized by the 2 disulfide bonds. One half of the structure is dominated by a 4 alpha-helix bundle with a left-handed twist; the helices are anti-parallel, with 2 overhand connections, which fall into a 2-stranded anti-parallel beta-sheet. The fourth alpha helix is important to the biological activity of the molecule. Interleukin-6 (IL-6) is an important proinflammatory and immunoregulatory cytokine expressed by various cells. Interleukin-6 has been shown to inhibit the growth of early stage and to promote the proliferation of advanced stage melanoma cells in vitro.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM HAc-NaAc, 150mM NaCl, pH 5.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : PVPPGEDSKDVAAPHRQPLTSSERIDKQIRYILDGIALRKETCNKSNMCESSKEALAENNLNLPKMAEKDGCFO
SGFNEETCLVKIITGLLEFEVYLEYLQNRFFESSEEQARAVQMSTKVLIQFLQKAKKNLDAITTPDPTTNASLLTKLQ
AQNQWLQDMTTHLILRSFKEFLQSSLRALRQM

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity : ED50 is less than 0.1 ng/ml. Specific Activity of 5.0 x 10⁷ IU/mg.