

32-7012: Recombinant Human Interleukin-2/IL-2

Gene : IL2
Gene ID : 3558
Uniprot ID : P60568

Description

Source: E.coli.
MW :15.4kD.

Recombinant Human Interleukin-2 is produced by our E.coli expression system and the target gene encoding Pro22-Thr153 is expressed. Recombinant Human Interleukin-2 is a highly purified protein with a molecular weight of approximately 15,300 Daltons. The chemical name is des-alanyl-1, serine-125 Human Interleukin-2. It is produced by recombinant DNA technology using a genetically engineered E. coli strain containing an analog of the human interleukin-2 gene. Genetic engineering techniques were used to modify the Human IL-2 gene, and the resulting expression clone encodes a modified Human IL-2. This recombinant form differs from native Interleukin-2 in following ways: 1) it is not glycosylated; 2) the molecule has no N-terminal alanine; 3) the molecule has serine substituted for cysteine at amino acid position 125; 4) the aggregation state of molecule is likely to be different from that of native IL-2.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM HAc-NaAc, 150mM NaCl, pH 4.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MPTSSSTKKTQLQLEHLLDLQMILNGINNYKNPKLTRMLTFKFYMPKKATELKHLCLEELKPLEEVNLQAQSKNFHLRPRDLISNINVIVLELKGSETTFMCEYADETATIVEFLNRWITFCQSIISTLT

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity : ED50 is less than 0.1 ng/ml. Specific Activity of 1.0 x 10⁷ IU/ mg.