

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7024: Recombinant Human Interleukin-17F/IL-17F

Gene ID: 112744 Uniprot ID: Q96PD4

Description

Source: E.coli. MW :30.1kD.

Recombinant Human Interleukin-17F is produced by our E.coli expression system and the target gene encoding Arg31-Gln163 is expressed. Interleukin-17 is a potent pro-inflammatory cytokine produced by activated memory T cells. There are at least six members of the IL-17 family in humans and in mice. Today, IL-17 represents a family of structurally related cytokines that share a highly conserved C-terminal region but differ from one another in their N-terminal regions and in their distinct biological roles. The six known members of this family, IL-17A through IL-17F, are secreted as homodimers. IL-17F also regulates cartilage matrix turnover and inhibits angiogenesis.

Product Info

Amount : 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MRKIPKVGHTFFQKPESCPPVPGGSMKLDIGIINENQRVSMSRNIESRSTSPWNYTVTWDPNRYPSEVVQAQCR

NLGCINAQGKEDISMNSVPIQQETLVVRRKHQGCSVSFQLEKVLVTVGCTCVTPVIHHVQ

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in 500mM Acetic Acid. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ \tilde{A} \square \hat{A} µg (1 IEU/ \tilde{A} \square \hat{A} µg) as determined by LAL test.

Biological Activity: ED50 is approximately 10 ng/ml. Specific Activity of 1.0 x 10^5 IU/mg