

32-7062: Recombinant Human Interleukin-37/IL-37

Gene : IL37
Gene ID : 27178
Uniprot ID : Q9NZH6

Description

Source: E.coli.
MW :18.7kD.

Recombinant Human Interleukin-37 is produced by our E.coli expression system and the target gene encoding Lys53-Asp218 is expressed. Human Interleukin family 1 Member 7 (IL1F7) is a member of the Interleukin 1 cytokine family. Five alternatively spliced transcript variants encoding distinct isoforms have been reported with distinct expression profiles. The longest IL1F7 transcript, referred to as IL1F7b or IL1F7 isoform 1, encodes a 218 amino acid residues proprotein containing a 45 amino acid propeptide, which is cleaved to generate mature protein. IL1F7b binds to IL18 Ra with low affinity but does not exert any IL18 agonistic or antagonistic effects. IL1F7b also binds interleukin 18 binding protein (IL-18BP), an inhibitory binding protein of interleukin 18 (IL-18), and subsequently forms a complex with IL18 receptor beta subunit, and through which it inhibits the activity of IL-18.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 2mM DTT, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MKNLNPKKFSIHDQDHKVLVLD SGNLI AVDPDKNYIRPEIFFALASSLSASA EKGSPILLGVSKGEFCLYCDKDKG QSHPSLQLKKEKLMKLA AQKESARRPFIFYRAQVGSWNMLESA AHPGWFICTSCNCNEPVGVTDFENRKHIE FSFQPVCKAEMSPSEVSD

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.