

32-7087: Recombinant Human beta-Arrestin 1/ARRB1 (C-6His)

Gene : ARRB1
Gene ID : 408
Uniprot ID : P49407

Description

Source: E.coli.
MW :48.13kD.

Recombinant Human beta-Arrestin 1 is produced by our E.coli expression system and the target gene encoding Met1-Arg418 is expressed with a 6His tag at the C-terminus. beta-Arrestin-1 (ARRB1) is a cytoplasmic protein that belongs to the arrestin family. ARRB1 is expressed at high levels in peripheral blood leukocytes and the central nervous system. ARRB1 regulates agonist-mediated G-protein coupled receptor (GPCR) signaling by mediating both receptor desensitization and resensitization processes. ARRB1 acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. ARRB1 is believed to play a major role in regulating receptor-mediated immune functions. ARRB1 is involved in Toll-like receptor and IL-1 receptor signaling through the interaction with TRAF6.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MGDKGTRVFKKASPNGLTVYLGKRDFVDHIDLVDVDPVGVVLDPEYLKERRVYVTLTCAFRYGGREDLDVLGLTFRKDLFVANVQSFPAPEDKKPLRLQERLIKKLGEHAYPFTFEIPPNLPCSVTLQPGPEDTGKACGV DYE VKAFCAENLEEKIHKRNSVRLVIRKVQYAPERPGPQPTAETTRQFLMSDKPLHLEASLDKEIYYHGEPISVNVHVTNNTNKT V K K I K I S V R Q Y A D I C L F N T A Q Y K C P V A M E E A D D T V A P S S T F C K V Y T L T P F L A N N R E K R G L A L D G K L K H E D T N L A S S T L L R E G A N R E I L G I I V S Y K V K V K L V V S R G G L L G D L A S S D V A V E L P F T L M H P K P K E E P P H R E V P E N E T P V D T N L I E L D T N D D D I V F E D F A R Q R L K G M K D D K E E E E D G T G S P Q L N N R L E H H H H H H

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.