

## 32-7105: Recombinant Human C-X-C Motif Chemokine 12/CXCL12/SDF-1 (22-93)

**Gene :** CXCL12

**Gene ID :** 6387

**Uniprot ID :** P48061

### Description

Source: E.coli.

MW :8.53kD.

Recombinant Human C-X-C Motif Chemokine 12 is produced by our E.coli expression system and the target gene encoding Lys22-Met93 is expressed. Stromal Cell-Derived Factor-1 (SDF-1) is a chemokine member of the intercrine family. SDF1 is expressed as five isoforms that differ only in the C terminal tail. SDF1a and SDF1 beta are identical except for the four residues present in the C-terminus of SDF1 beta but absent from SDF1a. SDF1 isoforms interact with CXCR4 and CXCR7 receptors on the cell surface, and can also bind syndecan4. SDF1 is known to influence lymphopoiesis, regulate patterning and cell number of neural progenitors, and promote angiogenesis. It also enhances the survival of myeloid progenitor cells.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** KPVLSYRCPCRFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVCIDPKLKWIQEYLEKALNKRFKM

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.