

32-7125: Recombinant Human Grancalcin/GCA (N-GST)

Gene : GCA
Gene ID : 25801
Uniprot ID : P28676

Description

Source: E.coli.
MW :50.3kD.

Recombinant Human Grancalcin is produced by our E.coli expression system and the target gene encoding Met1-Ile217 is expressed with a GST tag at the N-terminus. Grancalcin (GCA) is a member of the penta EF hand subfamily which includes sorcin, calpain and ALG2. Grancalcin is highly expressed bone marrow and also can detected in neutrophils and macrophages. Grancalcin interacts with L-plastin which known to have actin bundling activity. It indicates that Grancalcin may play an important role in the adhesion of neutrophils to fibronectin. Furthermore, Grancalcin localization is dependent upon calcium and magnesium. It associates with both the granule and membrane fractions, which suggested a role for grancalcin in granule-membrane fusion and degranulation.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl, 500mM NaCl, pH 8.5 .
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MSPILGYWKIKGLVQPTRLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPYYIDGDVKLTQSMAIRYIA
DKHNMLGGCPKERAISMLEGAVLDIRYGVSRAYSKDFETLKVDFLSKLPMLKMFEDRLCHKTYLNGDHVTH
PDFMLYDALDVVLYMDPMCLDAFPKLVCFKKRIEAIPIQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPR
GSMAYPGYGGGFGNFSIQVPGMQMGQVPETGPAILLDGYSGPAYSDTYSSAGDSVYTYFSAVAGQDGEVDA
EELQRCLTQSGINGTYSPFSLETCRMIAMLDRDHTGKMGFNAFKELWAALNAWKENFMTVDQDGSSTVEHH
ELRQAIGLMGYRLSPQTLTTIVKRYSKNGRIFDDYVACCVKLRALTDFFRKRDLHLQQGSANFIYDDFLQGTMAI

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.