

## 32-7174: Recombinant Human Myc-Associated Factor X/MAX (C-6His)

**Gene :** MAX  
**Gene ID :** 4149  
**Uniprot ID :** P61244

### Description

Source: E.coli.  
MW :18.27kD.

Recombinant Human Myc-Associated Factor X is produced by our E.coli expression system and the target gene encoding Met1-Ser151 is expressed with a 6His tag at the C-terminus. Myc-Associated Factor X (MAX) is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It contains 1 basic helix-loop-helix (bHLH) domain. It is found in the brain, heart, and lung at high levels while lower levels are seen in the liver, kidney, and skeletal muscle. MAX forms a sequence-specific DNA-binding protein complex with MYC or MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC-MAX complex is a transcriptional activator, whereas the MAD-MAX complex is a repressor. It may repress transcription via the recruitment of a chromatin remodeling complex containing H3 'Lys-9' histone methyltransferase activity.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl, 50mM Imidazole, 250mM NaCl, pH 8.5.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MSDNDDIEVESDADKRAHNNALERKRRDHKDSFHSRLRDSVPSLQGEKASRAQILDKATEYIQYMRRKNHHTHQ QDIDDLKRQNALLEQQVRALEKARSSAQLQTNYPSSDNSLYTNAKGSTISAFDGGSDSSSESEPEEPQSRKCLR MEASLEHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.