

## 32-7178: Recombinant Human EIF4E-Binding Protein 1/EIF4EBP1 (N-6His)

**Gene :** EIF4EBP1

**Gene ID :** 1978

**Uniprot ID :** Q13541

### Description

Source: E.coli.

MW :14.74kD.

Recombinant Human EIF4E-Binding Protein 1 is produced by our E.coli expression system and the target gene encoding Met1-Ile118 is expressed with a 6His tag at the N-terminus. Eukaryotic Translation Initiation Factor 4E-Binding Protein 1 (4EBP1) is a member of the eIF4E-binding protein family. 4EBP1 regulates eIF4E activity by preventing its assembly into the eIF4F complex. 4EBP1 mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase and mTORC1 pathways. Non-phosphorylated 4EBP1 competes with EIF4G1/EIF4G3 to interact with EIF4E. 4EBP1 is phosphorylated in response to various signals including insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation. 4EBP1 has a role in progression of breast neoplasms through cell signaling.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** MGSSHHHHHHSSGLVPRGSHMSGSSCSQTPSRAIPATRRVVLGDGVQLPPGDYSTTPGGTLFSTTPGGTRII  
YDRKFLMECRNSPVTKTPRDLPTIPGVTSPSSDEPPMEASQSHLRNSPEDKRAGGEESQFEMDI

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.