

32-7185: Recombinant Human Annexin A6/ANXA6 (C-6His)

Gene : ANXA6
Gene ID : 309
Uniprot ID : P08133

Description

Source: E.coli.
MW :76.9kD.

Recombinant Human Annexin A6 is produced by our E.coli expression system and the target gene encoding Ala2-Asp673 is expressed with a 6His tag at the C-terminus. Annexin A6 (ANXA6) belongs to a family of calcium-dependent membrane and phospholipid binding proteins. Annexin A6 is a secreted protein and locates on the cell surface. Annexin A6 contains 8 annexin repeats separated by linking sequences of variable lengths. A pair of annexin repeats may form one binding site for calcium and phospholipid. ANXA6 is involved in the regulation of the release of Ca²⁺ from intracellular stores and may be associated with CD21. In addition, ANXA6 has been implicated in mediating the endosome aggregation and vesicle fusion in secreting epithelia during exocytosis.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.
Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : AKPAQGAKYRGSIHDFPGFDPNQDAEALYTAMKGFSGDKEAILDIITSRSNRQRQEVCSYKSLYKGLDIADLKY
ELTGKFERLIVGLMRPPAYCDAKEIKDAISGIGTDEKCLIEILASRTNEQMHQLVAAYKDAYERDLEADIIGDTSG
HFQKMLVLLQGTREEDDVSEDLVQQDVQDLYEAGELKWTGDEAQFIYILGNRSKQHLRRLFVDFEYLKTTGKP
IEASIRGELSGDFEKLMLAVVKCIRSTPEYFAERLFKAMKGLGTRDNTLRIMVSRSELDMLDIREIFRTKYEKSLSY
MIKNDTSGEYKTKLLKSGGDDAAGQFFPEAAQVAYQMWELSAVARVELKGTVPANDFNPADAKALRKA
MKGLGTDEDTIIDIITHRSNVQRQQIRQTFKSHFGRDLMTDLKSEISGDLARLILGLMPPAHYDAKQLKKAMEG
AGTDEKALIEILATRTNAEIRAINAEYKEDYHKSLEDALSSDTSGHFRRILISLATGHREEGGENLDQAREDAQVA
AEILEIADTPSGDKTSLETRFMTILCTRSYPHLRRVQEFIKMTNYDVEHTIKKEMSGDVRDAFVAIVQSVKKNKPL
FFADKLYKSMKGAGTDEKTLTRIMVSRSEIDLNIRREFIEKYDKSLHQAIEGDTSGDFLKALLALCGGEDLEHHH
HHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.