

32-7197: Recombinant Human EIF1A, X-Chromosomal/EIF1AX (N-6His)(Discontinued)

Gene : EIF1AX
Gene ID : 107984923
Uniprot ID : P47813

Description

Source: E.coli.
MW :18.6kD.

Recombinant Human EIF1AX is produced by our E.coli expression system and the target gene encoding Met1-Ile144 is expressed with a 6His tag at the N-terminus. Eukaryotic Translation Initiation Factor 1A, X-Chromosomal (EIF1AX) is an essential eukaryotic translation initiation factor that belongs to the eIF-1A family. EIF1AX is required for the binding of the 43S complex (a 40S subunit, eIF2/GTP/Met-tRNAⁱ and eIF3) to the 5' end of capped RNA and has been shown to interact with IPO13. EIF1AX contains one S1-like domain and seems to be required for maximal rate of protein biosynthesis. Enhances ribosome dissociation into subunits and stabilizes the binding of the initiator Met-tRNA(I) to 40 S ribosomal subunits.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.
Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MGSSHHHHHHSSGLVPRGSHMPKNKGKGGKNNRRRGKNESEKRELVFKEDGQEYAQVIKMLGNRLEAM
CFDGVKRLCHIRGKLRKKVWINTSDIILVGLRDYQDNKADVILKYNADERSLKAYGELPEHAKINETDTFGPGD
DDEIQFDDIGDDEDEDIDDI

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.