

32-7250: Recombinant Human Apoptosis-Inducing Factor 1 Mitochondrial/AIFM1 (N-6His)

Gene : AIFM1
Gene ID : 9131
Uniprot ID : O95831

Description

Source: E.coli.
MW :56.2kD.

Recombinant Human Apoptosis-Inducing Factor 1 is produced by our E.coli expression system and the target gene encoding Glu121-Asp613 is expressed with a 6His tag at the N-terminus. Apoptosis-Inducing Factor 1, Mitochondrial (AIFM1) is a flavoprotein essential for nuclear disassembly in apoptotic cells that is found in the mitochondrial intermembrane space in healthy cells. During apoptosis, it is translocated from the mitochondria to the nucleus to function as a proapoptotic factor in a caspase-independent pathway, while in normal mitochondria, it functions as an antiapoptotic factor via its oxidoreductase activity. The soluble form (AIFsol) found in the nucleus induces parthanatos i.e., caspase-independent fragmentation of chromosomal DNA. AIFM1 interacts with EIF3G, and thereby inhibits the EIF3 machinery and protein synthesis, and activates caspase-7 to amplify apoptosis. It binds to DNA in a sequence-independent manner and plays a critical role in caspase-independent, pyknotic cell death in hydrogen peroxide-exposed cells.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MGSSHHHHHSSGLVPRGSHMEEVPQDKAPSHVFPFLIGGGTAAFAAARSIRARDPGARVLIVSEDPPELPYMR PPLSKELWFSDDPNVTKLRFKQWNGKERSIYFQPPSFYVSAQDLPHIENGGVAVLTGKKVVQLDVRDNMVKL NDGSQITYEKLIATGGTPRSLSAIDRAGAEVKSRTTLFRKIGDFRSLEKISREVKSITIIGGGFLGSELACALGRKA RALGTEVIQLFPEKGNMGKILPEYLSNWTMEKVRREGVKVMPNAIVQSVGVSSGKLLIKLDGRKVVTDHIVAA VGLEPNVELAKTGGLEIDSDFGGFRVNAELQARSNIWVAGDAACFYDIKLGRRRVEHHDHVVSGRLAGENM TGAAPYWHQSMFWSDLGPDVGYEAGLVDSLLPTVGVFAKATAQDNPKSATEQSGTGIRSESETESEASEITI PPSTPAVPQAPVQGEDYGKGVIFYLRDKVVVGVILWNIFNRMPIARKIIKDGEQHEDLNEVAKLFNIHED

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.