

36-1491: Monoclonal Antibody to MyoD1 (Rhabdomyosarcoma Marker)(5.8A + MYD712)(Discontinued)

Clonality :	Monoclonal
Clone Name :	5.8A + MYD712
Application :	FACS,IF,IHC
Reactivity :	Human,Mouse,Rat,Chicken
Gene :	MYOD1
Gene ID :	4654
Uniprot ID :	P15172
Format :	Purified
Alternative Name :	MYOD1,BHLHC1,MYF3,MYOD
Isotype :	Mouse IgG
Immunogen Information :	Recombinant mouse MyoD1 protein (5.8A); Recombinant human MyoD1 protein (MYD712)

Description


Recognizes a phosphor-protein of 45kDa, identified as MyoD1. This MAb does not cross react with myogenin, Myf5, or Myf6. Antibody to MyoD1 labels the nuclei of myoblasts in developing muscle tissues. MyoD1 is not detected in normal adult tissue, but is highly expressed in the tumor cell nuclei of rhabdomyosarcomas. Occasionally nuclear expression of MyoD1 is seen in ectomesenchymoma and a subset of Wilms tumors. Weak cytoplasmic staining is observed in several non-muscle tissues, including glandular epithelium and also in rhabdomyosarcomas, neuroblastomas, Ewing's sarcomas and alveolar soft part sarcomas.

Product Info

Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Flow Cytometry (0.5-1 µg/million cells in 0.1ml); Immunofluorescence (0.5-1 µg/ml); Immunohistology (Formalin-fixed) (0.5-1 µg/ml for 30 minutes at RT); (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes); Optimal dilution for a specific application should be determined.



Formalin-fixed, paraffin-embedded human Rhabdomyosarcoma stained with MyoD1 Monoclonal Antibody (5.8A + MYD712)