

32-7259: Recombinant Human SGTA (N-6His)

Gene : SGTA
Gene ID : 6449
Uniprot ID : O43765

Description

Source: E.coli.
MW :36.2kD.

Recombinant Human SGTA is produced by our E.coli expression system and the target gene encoding Met1-Glu313 is expressed with a 6His tag at the N-terminus. Small Glutamine-Rich Tetratricopeptide Repeat-Containing Protein a (SGTA) is an ubiquitously expressed protein which belongs to the SGT Family. SGTA contains three TPR Protein-Protein Interaction Duplicates. SGTA is a co-chaperone that binds directly to HSC70 and HSP70 and regulates their ATPase activity. SGTA is capable of interacting with the major nonstructural protein of Parvovirus H-1 and 70-kDa heat shock cognate protein. It interacts with NS1 from Parvovirus H-1, with Vpu and Gag from HIV-1. It also interacts with SARS-CoV Accessory Protein 7a, DNAJC5 and DNAJC5B. However, its function is not known. Since this transcript is expressed ubiquitously in various tissues, SGTA may serve a housekeeping function.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 150mM NaCl, pH 8.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MGSSHHHHHSSGLVPRGSHMDNKKRLAYAIQFLHDQLRHGGLSSDAQESLEVAIQCLEAFGVTVEDSDLA LPQTLPEIFEAAATGKEMPQDLRSPARTPPSEEDSAEAERLKTEGNEQMKVENFEAAVHFYGGKAIELNPNANAVYF CNRAAAYSCLGNYAGAVQDCERAICIDPAYSKAYGRMGLALSSLNKHVEAVAYYKKALELDPDNETYKSNLKIA ELKLRAPSPPTGGVGSFDIAGLLNPNPGFMSMASNLMNNPQIQQLMSGMISGGNNPLGTPGTSPSQNDLASLIQ AGQQFAQQMQQNPELIEQLRSQIRSRTPSASNDQEQE

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.