

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7278: Recombinant Human Tryptophan--tRNA Ligase, Cytoplasmic/WARS/TrpRS (N-6His)

Gene ID: 7453
Uniprot ID: P23381

## **Description**

Source: E. coli. MW :55.3kD.

Recombinant Human TrpRS is produced by our E.coli expression system and the target gene encoding Met1-Gln471 is expressed with a 6His tag at the N-terminus. There exists two types of tryptophanyl tRNA synthetases, the cytoplasmic form called WARS, the mitochondrial form called WARS2. WARS catalyzes the aminoacylation of tRNA (trp) with tryptophan and is induced by interferon. WARS regulates ERK, Akt, eNOS activation pathway, which are related with angiogenesis, cytoskelatal reorganization and shear stess-reponsive gene expression.

## **Product Info**

Amount:  $10 \mu g / 50 \mu g$ 

**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 150mM NaCl, pH 8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MGSSHHHHHHSSGLVPRGSHMPNSEPASLLELFNSIATQGELVRSLKAGNASKDEIDSAVKMLVSLKMSYKAA

AGEDYKADCPPGNPAPTSNHGPDATEAEEDFVDPWTVQTSSAKGIDYDKLIVRFGSSKIDKELINRIERATGQR PHHFLRRGIFFSHRDMNQVLDAYENKKPFYLYTGRGPSSEAMHVGHLIPFIFTKWLQDVFNVPLVIQMTDDEKY LWKDLTLDQAYSYAVENAKDIIACGFDINKTFIFSDLDYMGMSSGFYKNVVKIQKHVTFNQVKGIFGFTDSDCIG KISFPAIQAAPSFSNSFPQIFRDRTDIQCLIPCAIDQDPYFRMTRDVAPRIGYPKPALLHSTFFPALQGAQTKMSAS DPNSSIFLTDTAKQIKTKVNKHAFSGGRDTIEEHRQFGGNCDVDVSFMYLTFFLEDDDKLEQIRKDYTSGAMLT

GELKKALIEVLQPLIAEHQARRKEVTDEIVKEFMTPRKLSFDFQ

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.