

32-7280: Recombinant Human Activin Receptor 2B/Activin RIIB/ACVR2B (C-6His)

 Gene :
 ACVR2B

 Gene ID :
 93

 Uniprot ID :
 Q13705

Description

Source: Human Cells.

MW :14.37kD.

Recombinant Human Activin Receptor IIB is produced by our Mammalian expression system and the target gene encoding Ser19-Thr134 is expressed with a 6His tag at the C-terminus. Activin proteins that belong to the transforming growth factorbeta (TGF- beta) superfamily, exert their biological actions by binding to heteromeric receptor complexes of type I and type II serine/threonine kinase receptors. On ligand binding, type I and II receptors form a stable complex, resulting in phosphorylation of type I receptors by type II receptors with constitutive kinase activity, and subsequently initiates the activation of downstream molecules including the endogenous Smads. ActRIIB, also known as ActRIIB, is a type II receptor containing an extracellular domain (ECD), a transmembrane segment, and a cytoplasmic region that includes the kinase domain. ActRIIB is a receptor for activin A, activin B and inhibin A. Multiple ActRIIB isoforms can also be generated, which bind activin isoforms with different affinities.

Product Info

| Amount : Content : | 10 µg / 50 µg Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. |
|-----------------------|--|
| Storage condition : | Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months. |
| Amino Acid : | SGRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSGTIELVKKGCWLDDFNCYDRQEC VATEENPQVYFCCCEGNFCNERFTHLPEAGGPEVTYEPPPTAPTVDHHHHHH |

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 \tilde{A} $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ \tilde{A}] \hat{A} µg (1 IEU/ \tilde{A}] \hat{A} µg) as determined by LAL test.