

32-7282: Recombinant Human C1qTNF1/CTRP1 (C-6His)

Gene : C1QTNF1

Gene ID : 114897

Uniprot ID : Q9BXJ1

Description

Source: Human Cells.

MW :30.24kD.

Recombinant Human C1qTNF1 is produced by our Mammalian expression system and the target gene encoding Arg26-Pro281 is expressed with a 6His tag at the C-terminus. C1QTNF1 is a secreted protein, contains 1 C1q domain and 1 collagen-like domain. C1qTNF proteins constitute a highly conserved family of Acrp30/Adiponectin paralogs that share a modular organization comprising an N-terminal signal peptide, a short variable region, a collagenous domain and a C-terminal globular domain. C1qTNF proteins are predicted to have trimeric structures that assemble into hexameric and higher order molecular forms. C1QTNF1 is a novel adipokine, providing a significant framework to further address the physiological functions and mechanisms of the action of this family of secreted glycoproteins in normal and disease states. C1QTNF1 increases the production of aldosterone. C1QTNF1 is vastly expressed in obese subjects as well as up-regulated in hypertensive patients, C1QTNF1 is identified molecular link between obesity and hypertension. C1QTNF1 expression may be associated with a low-grade chronic inflammation status in adipose tissues.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : RVPHVQGEQQEWEGTEELPSPDHAERAEQHEKYRPSQDQGLPASRCLRCCDPGTSMPATAVPQINITLK
GEKGDRGDRGLQGKYGKTGSAGARGHTGPKGQKSMGAPGERCKSHYAAFSVGRKKPMHSHYYQTVIFDT
EFVNLVDHFNMTGKFCYVPGLYFFSLNVHTWNQKETYLHIMKNEEEVILFAQVGDERSIMQSQSLMLELREQ
DQVWVRLYKGERENAIFSEELDTYITFSGYLVKHATEPVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.