

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7306: Recombinant Human Coxsackievirus and Adenovirus Receptor/CAR/CXADR (C-6His)(Discontinued)

Gene ID: CXADR
Gene ID: 1525
Uniprot ID: P78310

Description

Source: Human Cells. MW :25.08kD.

Recombinant Human Coxsackievirus and Adenovirus Receptor is produced by our Mammalian expression system and the target gene encoding Leu20-Gly237 is expressed with a 6His tag at the C-terminus. Coxsackievirus and Adenovirus Receptor (CAR) belongs to the CTX family of the Ig superfamily. CXADR is a type I transmembrane glycoprotein and expressed in pancreas, brain, heart, small intestine, testis, prostate. It is a receptor that mediates gene transfer and also act as an adhesion molecule within junctional complexes, notably between epithelial cells lining body cavities and within myocardial intercalated discs. CXADR contains an extracellular domain, a transmembrane helix and a C-terminal intracellular domain. The C-terminal interacts with few cytoplasmic junctional proteins, microtubules and the actin cytoskeleton.

Product Info

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: LSITTPEEMIEKAKGETAYLPCKFTLSPEDQGPLDIEWLISPADNQKVDQVIILYSGDKIYDDYYPDLKGRVHFTSN

DLKSGDASINVTNLQLSDIGTYQCKVKKAPGVANKKIHLVVLVKPSGARCYVDGSEEIGSDFKIKCEPKEGSLPL QYEWQKLSDSQKMPTSWLAEMTSSVISVKNASSEYSGTYSCTVRNRVGSDQCLLRLNVVPPSNKAGVDHHHH

HH

Application Note

Storage condition:

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.