

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7332: Recombinant Human Kallikrein 4/KLK4 (C-6His)

**Gene :** KLK4 **Gene ID :** 9622 **Uniprot ID :** 09Y5K2

## **Description**

Source: Human Cells. MW:25.44kD.

Recombinant Human Kallikrein 4 is produced by our Mammalian expression system and the target gene encoding Ser27-Ser254 is expressed with a 6His tag at the C-terminus. Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many Kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen members of the Kallikrein subfamily located in a cluster on chromosome 19. Its encoded protein is secreted and may play a role in suppression of tumorigenesis in breast and prostate cancers. Alternate splicing of this gene results in multiple transcript variants encoding the same protein.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl, 150mM NaCl, pH 8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: SCSQIINGEDCSPHSQPWQAALVMENELFCSGVLVHPQWVLSAAHCFQNSYTIGLGLHSLEADQEPGSQMVE

ASLSVRHPEYNRPLLANDLMLIKLDESVSESDTIRSISIASQCPTAGNSCLVSGWGLLANGRMPTVLQCVNVSVV SEEVCSKLYDPLYHPSMFCAGGGQDQKDSCNGDSGGPLICNGYLQGLVSFGKAPCGQVGVPGVYTNLCKFTE

**WIEKTVQASVDHHHHHH** 

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.