

32-7353: Recombinant Human Osteonectin/SPARC/BM-40 (C-6His)(Discontinued)

Gene : SPARC
Gene ID : 6678
Uniprot ID : P09486

Description

Source: Human Cells.
MW :33.73kD.

Recombinant Human Osteonectin is produced by our Mammalian expression system and the target gene encoding Ala18-Ile303 is expressed with a 6His tag at the C-terminus. Secreted Protein Acidic and Rich in Cysteine (SPARC) is a secreted, evolutionarily conserved collagen-binding glycoprotein and belongs to the SPARC family. SPARC has 286 amino acids and contains an EF-hand in C-termina domain, a follistatin-like domain with Kazal-like sequences. There are two calcium binding sites, one binds 5 - 8 Ca²⁺ with a low affinity and other on an EF-hand loop that binds a Ca²⁺ ion with a high affinity. It is highly expressed in tissues undergoing morphogenesis, remodeling and wound repair. SPARC regulate cell growth through interactions with the extracellular matrix (ECM) and cytokines. SPARC bind to numerous proteins of the ECM, affect ECM protein expression, influence cellular adhesion and migration, and modulate growth factor-induced cell proliferation and angiogenesis. SPARC also binds several types of collagen, albumin, thrombospondin, PDGF and cell membranes.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : APQQEALPDETEVVEETVAEVTEVSVGANPVQVEVGEFDDGAEETEEVVAENPCQNHHCCKHGKVCELDENNTPMCVCQDPTSCPAPIGEFEKVCNSDNKTFDSSCHFFATKCTLEGTKKGHKLHLDYIGPCKYIPPLDSELTEFPLRMRDWLKNVLVTLYERDEDNLLTEKQKLRVKKIHENEKRLEAGDHPVELLARDFEKNYMYIFPVHWQFGQLDQHPIDGYLSHTELAPLRAPLIPMEHCTTRFFETCDLDNDKYIALDEWAGCFGIKQKDKDLVIVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.