

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7354: Recombinant Human Low-Density Lipoprotein Receptor Related Protein 12/LRP12 (C-6His)(Discontinued)

Gene ID: 29967 **Uniprot ID:** Q9Y561

Description

Source: Human Cells. MW :52.48kD.

Recombinant Human LRP12 is produced by our Mammalian expression system and the target gene encoding Asn28-Ile488 is expressed with a 6His tag at the C-terminus. Low-Density Lipoprotein Receptor-Related Protein 12 (LRP12) belongs to the LDLR family. LRP12 is a type I transmembrane protein annd widely expressed in heart, skeletal muscle, brain, lung, placenta and pancreas. LRP12 contains 2 CUB domain and 5 LDL-receptor class A domain. LRP12 has been shown to interact with GNB2L1, ZFYVE9 and ITGB1BP3. LRP12 is a receptor probably, which may be involved in the internalization of lipophilic molecules and/or signal transduction. In addition, LRP12 may act as a tumor suppressor.

Product Info

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20 $^{\circ}\text{C},$ though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: NGALAEHSENVHISGVSTACGETPEQIRAPSGIITSPGWPSEYPAKINCSWFIRANPGEIITISFQDFDIQGSRRCN

LDWLTIETYKNIESYRACGSTIPPPYISSQDHIWIRFHSDDNISRKGFRLAYFSGKSEEPNCACDQFRCGNGKCIP EAWKCNNMDECGDSSDEEICAKEANPPTAAAFQPCAYNQFQCLSRFTKVYTCLPESLKCDGNIDCLDLGDEID CDVPTCGQWLKYFYGTFNSPNYPDFYPPGSNCTWLIDTGDHRKVILRFTDFKLDGTGYGDYVKIYDGLEENPHK LLRVLTAFDSHAPLTVVSSSGQIRVHFCADKVNAARGFNATYQVDGFCLPWEIPCGGNWGCYTEQQRCDGYW HCPNGRDETNCTMCQKEEFPCSRNGVCYPRSDRCNYQNHCPNGSDEKNCFFCQPGNFHCKNNRCVFESWV

CDSQDDCGDGSDEENCPVIVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than $0.1 \text{ ng}/\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.