9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-7361: Recombinant Human Wnt Inhibitory Factor 1/WIF-1 (C-6His)

Gene : WIF1
Gene ID : 11197
Uniprot ID : Q9Y5W5

Description

Source: Human Cells.
MW :39.47kD.
Recombinant Human Wnt Inhibitory Factor 1 is produced by our Mammalian expression system and the target gene encoding Gly29-Trp379 is expressed with a 6His tag at the C-terminus. Wnt Inhibitory Factor 1 (WIF1) is a secreted protein, which binds WNT proteins and inhibits their activities. WNT proteins are extracellular signaling molecules involved in the control of embryonic development. WIF1 contains a WNT inhibitory factor (WIF) domain and 5 epidermal growth factor (EGF)-like domains. is found to be present in fish, amphibia and mammals. WIF1 is a recurrent target in human salivary gland oncogenesis.WIF1 may be involved in mesoderm segmentation. WIF1 is a tumor suppressor, specifically in nonfunctioning pituitary tumors.

Product Info

Amount :	10 μg / 50 μg
Content :	Lyophilized from a 0.2 μm filtered solution of 10mM HAc-NaAc, 150mM NaCl, 0.5% CHAPS, pH 4.0.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	GPPQEESLYLWIDAHQARVLIGFEEDILIVSEGKMAPFTHDFRKAQQRMPAIPVNIHSMNFTWQAAGQAEYFYE FLSLRSLDKGIMADPTVNVPLLGTVPHKASVVQVGFPCLGKQDGVAAFEVDVIVMNSEGNTILKTPQNAIFFKTC QQAECPGGCRNGGFCNERRICECPDGFHGPHCEKALCTPRCMNGGLCVTPGFCICPPGFYGVNCDKANCSTT CFNGGTCFYPGKCICPPGLEGEQCEISKCPQPCRNGGKCIGKSKCKCSKGYQGDLCSKPVCEPGCGAHGTCHE PNKCQCQEGWHGRHCNKRYEASLIHALRPAGAQLRQHTPSLKKAEERRDPPESNYIWVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 ÃÂμg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Endotoxin : Less than 0.1 ng/ÃÂμg (1 IEU/ÃÂμg) as determined by LAL test.