

## 32-7371: Recombinant Human Dickkopf-Related Protein 3/DKK3 (C-6His)(Discontinued)

**Gene :** DKK3  
**Gene ID :** 27122  
**Uniprot ID :** Q9UBP4

### Description

Source: Human Cells.  
MW :37.22kD.

Recombinant Human Dickkopf-Related Protein 3 is produced by our Mammalian expression system and the target gene encoding Ala22-Ile350 is expressed with a 6His tag at the C-terminus. Dickkopf-related protein 3 (DKK3) belongs to the DKK protein family including Dkk-1, 2, 3 and -4. DKK3 is a 350 amino acid secreted glycoprotein which is comprised of an N-terminal signal peptide and 2 conserved cysteine-rich domains that are separated by a 12 amino acid linker region. Dkk-3 also have one prokineticin domain. DKK3 is involved in embryonic development through its inhibition of the WNT signaling pathway. The Dkk family also includes Soggy, which is homologous to Dkk-3 but not to the other family members. Soggy has not been shown to inhibit Wnt signaling, and its role in the pathway is unclear.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** APAPTATSAPVKPGPALSYPQEEATLNEMFREVEELMEDTQHKLRSVEEMEAEAAAASSEVNLANLPSPYH  
NETNTDTKVGNNTIHVHREIHKITNNQTGQMVFSETVITSVGDEEGRRSHECIIDEDCGPSMYCQFASFQYTCQ  
PCRGQRMLCTRDSECCGDQLCVWGHCTKMATRGSNGTICDNQRDCQPGLCCAFQRGLLFPVCTPLPVEGEL  
CHDPASRLDLITWELEPDGALDRCPASGLLCQPHSHSLVYVCKPTFVGSRDQDGEILLPREVPDEYEVGSFM  
EEVRQELEDLERSLTEEMALGEPAAAAAALLGGEEIVDHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.