

## 32-7378: Recombinant Human AGER/RAGE (C-6His)

**Gene :** AGER  
**Gene ID :** 177  
**Uniprot ID :** Q15109

### Description

Source: Human Cells.  
MW :35.2kD.

Recombinant Human AGER is produced by our Mammalian expression system and the target gene encoding Ala23-Ala344 is expressed with a 6His tag at the C-terminus. Advanced Glycosylation End Product-Specific Receptor (AGER) belongs to the immunoglobulin superfamily of cell surface molecules. It lies within the major histocompatibility complex (MHC) class III region on chromosome 6. Besides AGEs, AGER is also able to bind other ligands which is thought to result in pro-inflammatory gene activation. It is known that AGER serve as a mediator of both acute and chronic vascular inflammation in certain conditions such as atherosclerosis and in particular as a complication of diabetes. Furthermore, it plays an important role in regulating the production/expression of TNF-alpha, oxidative stress, and endothelial dysfunction in type 2 diabetes.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** AQNITARIGEPLVLKCKGAPKPPQRLEWKLNTGRTEAWKVLSPQGGGPWDSVARVLPNGSLFLPAVGIQDEGI  
FRCQAMNRNGKETKSNYRVRVYQIPGKPEIVDSASELTAGVPNKVGTVCVSEGSYPAGTLSWHLDGKPLVPNEK  
GVSVKEQTRRRHPETGLFTLQSELMVTPARGGDRPRTFSCSFSPGLPRHRALRTAPIQPRVWEPVPLEEVQLVVE  
PEGGAVAPGGTVTLTCEVPAQPSPQIHWKMDGVPLPLPPSPVLILPEIGPQDQGTYSVATHSSHGPPQESRAVS  
ISIIPEGGEGPTAGSVGGSGGLGTLALAVDHHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.