

32-7384: Recombinant Human Azurocidin/CAP37/HBP (C-6His)

Gene : AZU1
Gene ID : 566
Uniprot ID : P20160

Description

Source: Human Cells.
MW :25.24kD.

Recombinant Human Azurocidin is produced by our Mammalian expression system and the target gene encoding Ile27-Pro250 is expressed with a 6His tag at the C-terminus. Azurocidin is an Azurophil granule antibiotic protein, with monocyte chemotactic and antibacterial activity. The Azurophil granules, specialized lysosomes of the neutrophil, contain at least 10 proteins implicated in the killing of microorganisms. Azurocidin is a member of the serine protease family that includes Cathepsin G, Neutrophil Elastase (NE), and Proteinase 3 (PR3), however, Azurocidin is not a serine proteinase since the active site serine and histidine residues are replaced. Human Azurocidin together with NE and PR3 are expressed coordinately and are packaged together into azurophil granules during neutrophil differentiation. Azurocidin has been identified as a modulator of endothelial permeability and an important multifunctional inflammatory mediator. Neutrophils arriving first at sites of inflammation release Azurocidin which acts in a paracrine fashion on endothelial cells causing the development of intercellular gaps and allowing leukocyte extravasation. Azurocidin thus be regarded as a reasonable therapeutic target for a variety of inflammatory disease conditions.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM HEPES, 150mM NaCl, pH 7.5.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : IVGGRKARPRQFPFLASIQNQGRHFCGGALIHARFVMTAASCFQSQNPGVSTVVLGAYDLRRRERQSRQTFSIS
SMSSENGYDPQQNLNDLMLLQLDREANLTSSVTILPLPLQ NATVEAGTRCQVAGWGSQRSGGRLSRFPFVN
TVTPEDQCRPNNVCTGVLTRRGGICNGDGGTPLVCEGLAHGVASFSLGPCGRGPDFFTRVALFRDWIDGVLN
NPGPGPVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.