

32-7394: Recombinant Human Poliovirus Receptor-Related Protein 2/PVRL2/CD112 (C-6His)

Gene : NECTIN2
Gene ID : 5819
Uniprot ID : Q92692

Description

Source: Human Cells.
MW :36.58kD.

Recombinant Human Nectin-2 is produced by our Mammalian expression system and the target gene encoding Gln32-Leu360 is expressed with a 6His tag at the C-terminus. CD112 is a type I transmembrane glycoprotein belonging to the Immunoglobulin superfamily. It comprises one Ig-like V-type domain and two Ig-like C2-type domains in the extracellular region. The V domain is believed to mediate nectin binding to its ligands. Nectin2 is known to bind the pseudorabies virus, and herpes simplex virus2 (HSV2), involving in cell to cell spreading of these viruses. It does not bind poliovirus. As a homophilic adhesion molecule, CD112 is found concentrated in adherens junctions, and exists on neurons, endothelial cells, epithelial cells and fibroblasts. CD112 has been identified as the ligand for DNAM-1 (CD226), and the interaction of CD226/CD112 mediates cytotoxicity and cytokine secretion by T and NK cells. The costimulatory responses may be a critical component in allergic reactions and may therefore become targets for anti-allergic therapy.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : QDVRVQVLPEVRGQLGGTVLPCHELLPPVPGLYISLVTWQRPDAPANHQNVAAAFHPKMGPSFSPKPGSERLS FVSAKQSTGQDTEAELQDATALALHGLTVEDEGNYTCEFATFPKGSVRGBMTWLRVIAKPKNQAEAKVTFSQDP TTVALCISKEGRPPARISWLSSLDWEAKETQVSGTLAGTVTVTSRFTLVPSGRADGVTVTCKVEHESFEEPALIP VTLSVRYPPEVSISGYDDNWYLGRTDATALSCDVRSNPEPTGYDWSTTSGTFPTSAVAQGSQQLVIHAVDSLNTT FVCTVTNAVGMGRAEQVIFVRETPRASPRDVGPLVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.