

32-7415: Recombinant Human C-X3-C Motif Chemokine 1/CX3CL1/Fractalkine (C-6His)

Gene : CX3CL1

Gene ID : 6376

Uniprot ID : P78423

Description

Source: Human Cells.

MW :34.36kD.

Recombinant Human C-X3-C Motif Chemokine 1 is produced by our Mammalian expression system and the target gene encoding Gln25-Arg339 is expressed with a 6His tag at the C-terminus. Human Fractalkine (CX3CL1) is a member of the CX3C family of chemokines. Human Fractalkine contains both chemokine and mucin domain. The soluble form of Fractalkine is chemotactic for T-cells and monocytes, but not for neutrophils. The membrane bound form of Fractalkine promotes leukocytes adhesion to endothelial cells. Fractalkine regulates leukocyte adhesion and migration processes at the endothelium and binds to CX3CR1. Natural Human Fractalkine is produced as a long protein (373-amino acid). The mucin-like stalk permits it to bind to the cell surface.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : QHHGVTKCNITCSKMTSKIPVALLIHYQQNQASCGRKRAIILETRQHRLFCADPKQWVKDAMQHLDRQAAALT
RNGGTFEKQIGEVKPRTPAAGGMDESVVLEPEATGESSLEPTSSQEAQRALGTSPQLPTGVTGSSGTRLPP
TPKAQDGGPVGTELFVRVPPVSTAATWQSSAPHQPGPSLWAEAKTSEAPSTQDPSTQASTASSPAPEENAPSEG
QRVWGGQSPRPENSLEREEMGPVPAHTDAFQDWGPGSMAHVSVPVSSEGTPSREPVASGSWTPKAEPI
HATMDPQRLGVLITPVPDAQAATRVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.