

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7416: Recombinant Human Cystatin SN/CST1 (C-6His)

**Gene ID:** 1469 **Uniprot ID:** P01037

## **Description**

Source: Human Cells. MW:15.35kD.

Recombinant Human Cystatin SN is produced by our Mammalian expression system and the target gene encoding Trp21-Ser141 is expressed with a 6His tag at the C-terminus. Cystatin-SN is a member of family 2 of the Cystatin superfamily and a cysteine proteinase inhibitor found in saliva, tears, urine, and seminal fluid. Together with Cystatins S and SA, it is produced by the salivary gland and secreted largely in the submandibular/sublingual saliva. Cystatin-SN inhibits members of the Papain family including Cathepsins B, C, H and L.

## **Product Info**

Amount:  $10 \mu g / 50 \mu g$ 

**Content :** Lyophilized from a 0.2 μm filtered solution of 20mM MES, 150mM NaCl, pH 5.5.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: WSPKEEDRIIPGGIYNADLNDEWVQRALHFAISEYNKATKDDYYRRPLRVLRARQQTVGGVNYFFDVEVGRTIC

TKSQPNLDTCAFHEQPELQKKQLCSFEIYEVPWENRRSLVKSRCQESVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.