

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7440: Recombinant Human Ly6/PLAUR Domain-Containing Protein 3/LYPD3/C4.4A (C-6His)

Gene ID: 27076 **Uniprot ID:** 095274

Description

Source: Human Cells. MW:27.89kD.

Recombinant Human LYPD3 is produced by our Mammalian expression system and the target gene encoding Leu31-His286 is expressed with a 6His tag at the C-terminus. Ly6/PLAUR domain containing3 (LYPD-3) is a GPI-linked protein. The structure of LYPD-3 is similar to the urokinasetype plasminogen activator receptor (uPAR). LYPD-3 is a 6 -100 kDa molecule with variable cell type-specific N-O-linked glycosylation, mature human LYPD-3 contains two uPAR/Ly6 domains and a Ser/Thr/Prorich (STP) region includes a protease sensitive site . The interaction of LYPD-3 with Laminin 1 and 5 on neighboring cells promotes the adhesion, spreading, and migration of tumor cells. LYPD-3 additionally interacts with Galectin-3 and the anterior gradient proteins AG-2 and AG-3. LYPD-3 overexpression in non-small cell lung cancer is predictive of increased mortality.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid: LECYSCVQKADDGCSPNKMKTVKCAPGVDVCTEAVGAVETIHGQFSLAVRGCGSGLPGKNDRGLDLHGLLAFI

QLQQCAQDRCNAKLNLTSRALDPAGNESAYPPNGVECYSCVGLSREACQGTSPPVVSCYNASDHVYKGCFDG NVTLTAANVTVSLPVRGCVQDEFCTRDGVTGPGFTLSGSCCQGSRCNSDLRNKTYFSPRIPPLVRLPPPEPTTV

ASTTSVTTSTSAPVRPTSTTKPMPAPTSQTPRQGVEHVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.