

32-7475: Recombinant Human Oxidized Low-Density Lipoprotein Receptor 1/OLR1 (C-6His)

Gene : OLR1
Gene ID : 4973
Uniprot ID : P78380

Description

Source: Human Cells.
MW :25.39kD.

Recombinant Human LOX-1 is produced by our Mammalian expression system and the target gene encoding Ser61-Gln273 is expressed with a 6His tag at the C-terminus. Oxidized Low-Density Lipoprotein Receptor 1 (Ox-LDL Receptor 1) is a secreted, single-pass type II membrane protein which belongs to the C-type lectin superfamily. Ox-LDL Receptor 1 is expressed at high levels in endothelial cells and vascular-rich organs such as placenta, lung, liver, brain, aortic intima, bone marrow, spinal cord and substantia nigra. The expression of Ox-LDL Receptor 1 is induced by inflammatory cytokines such as TNF, IFNG and IL6 by pathological conditions, such as hyperlipidemia, hypertension and diabetes mellitus. Ox-LDL Receptor 1 mediates the recognition, internalization and degradation of oxidatively modified low density lipoprotein (OxLDL) by vascular endothelial cells. Ox-LDL Receptor 1 association with oxLDL induces the activation of NF-kappa-B through an increased production of intracellular reactive oxygen and a variety of pro-atherogenic cellular responses including a reduction of nitric oxide (NO) release, monocyte adhesion and apoptosis. Ox-LDL Receptor 1 also binds to oxLDL, which acts as a receptor for the HSP70 protein involved in antigen cross-presentation to naive T-cells in dendritic cells, thereby participating in cell-mediated antigen cross-presentation. It also participates in inflammatory process, by acting as a leukocyte-adhesion molecule at the vascular interface in endotoxin-induced inflammation.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : SQVSDLLTQEQANLTHQKKKLEGQISARQQAEASQSENELKEMIETLARKLNEKSKEQMELHHQNLNLQET
LKRVANCSAPCPQDWIWHGENCYLFSSGSFNWEKSQEKCLSLDAKLLKINSTADLDFIQQAISYSSFPFWMGLS
RRNPSYPWLWEDGSPLMPHLFRVRGAVSQTYPSGTCAIYIRGAVYAENCILAAFSICQKKANLRAQVDHHHHH
H

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.