

32-7479: Recombinant Human Podoplanin/PDPN/Aggrus (C-6His)

Gene : PDPN
Gene ID : 10630
Uniprot ID : Q86YL7

Description

Source: Human Cells.
MW :12.16kD.

Recombinant Human Podoplanin is produced by our Mammalian expression system and the target gene encoding Ala23-Leu131 is expressed with a 6His tag at the C-terminus. Podoplanin is a type-1 transmembrane protein that belongs to Podoplanin family. PDPN expressed in various specialized cell types throughout the body. It highly expressed in placenta, lung, skeletal muscle and brain, weakly expressed in brain, kidney and liver. In placenta, PDPN expressed on the apical plasma membrane of endothelium, in lung, expressed in alveolar epithelium. PDPN physiological function is related to its mucin-type character. PDPN may be involved in cell migration and/or actin cytoskeleton organization. When expressed in keratinocytes, induces changes in cell morphology with transfected cells showing an elongated shape, numerous membrane protrusions, and major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion. It requires for normal lung cell proliferation and alveolus formation at birth and Induces platelet aggregation. Nevertheless, it doesn't have any effect on amino acid transport and the aquaporin-type water channels.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : ASTGQPEDDTETTGLEGGVAMPGAEDDVVTPGTSSEDRYKSLTTLVATSVNSVTGIRIEDLPTSESTVHAQEQS
PSATASNVATSHSTEKVDGDTQTTVEKDGLSTVTLVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.