

32-7480: Recombinant Human Plexin Domain-Containing Protein 1/PLXDC1 (C-6His)

Gene : PLXDC1

Gene ID : 57125

Uniprot ID : Q8IUU5

Description

Source: Human Cells.

MW :46.53kD.

Recombinant Human Plexin Domain-Containing Protein 1 is produced by our Mammalian expression system and the target gene encoding Leu19-Thr426 is expressed with a 6His tag at the C-terminus. Plexin Domain-Containing Protein 1 (PLXDC1) is a single-pass type I membrane protein that belongs to the plexin family. Secreted PLXDC1 is localized predominantly at the tight junctions of vascular endothelial cells and to a lesser extent at the luminal surface of vascular endothelial cells. PLXDC1 is expressed in fibrovascular membrane with increased expression in individuals with proliferative diabetic retinopathy. It can detect in endothelial cells from colorectal cancer, and in endothelial cells from primary cancers of the lung, liver, pancreas, breast and brain. PLXDC1 interacts with NID1 and may also interact with CTTN. It plays a important role in endothelial cell capillary morphogenesis, the proliferation and maintenance of neovascular endothelial cells in the formation of fibrovascular membranes (FVMs).

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 5% Threhalose, pH 7.2.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : LSPQPGAGHDEGPGSGWAAKGTVRGWNRRARESPPGHVSEPDRTQLSQDLGGGTLAMDTLPDNRTRVVEDN
HSYYVSRLYGPSEPHSRELWVDVAEANRSQVKIHTILSNTHRQASRVVLSFDFPFYGHPLRQITMATGGFIFMG
DVIHRMLTATQYVAPLMANFNPGYSDNSTVVYFDNGTVFVVQWDHVYLQGWEDKGSFTFQAALHHDGRIVF
AYKEIPMSVPEISSQHPVKTGLSDAFAMILNPSDPVPESRRRSIFEYHRIELDPSKVTSMSAVEFTPLPTCLQHRSC
DACMSSDLTFNCSWCHVLQRCSSGFDRYRQEWMDYGCAQEAEGRMCEDFQDEHDHSASPDTSFSPYDGD
TTSSSLFIDSLTTEDDTKLNPNYAGGDGLQNNLSPKTKGTPVHLGTVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.