

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7507: Recombinant Human Apolipoprotein M/ApoM (C-6His)

Gene ID: 55937 **Uniprot ID:** 095445

Description

Source: Human Cells. MW:22.29kD.

Recombinant Human Apolipoprotein M is produced by our Mammalian expression system and the target gene encoding Met1-Asn188 is expressed with a 6His tag at the C-terminus. Apolipoprotein M is a secreted protein which belongs to the Lipocalin family. ApoM often presents in high density lipoprotein (HDL) and to a lesser extent in triglyceride-rich lipoproteins (TGRLP) and low density lipoproteins (LDL). The ApoM gene encoded protein is expressed in liver and kidney, secreted through the plasma membrane but remains membrane-bound. ApoM probably involved in lipid transport. ApoM can bind sphingosine-1-phosphate, myristic acid, palmitic acid and stearic acid, retinol, all-trans-retinoic acid and 9-cis-retinoic acid. The expression of ApoM could be regulated by platelet activating factor (PAF), Transforming Growth Factors (TGF), Insulin-Like Growth factor (IGF) and Leptin.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

Storage condition : Reconstituted protein solution can be stored a samples are stable at -20°C for 3 months.

Amino Acid: MFHQIWAALLYFYGIILNSIYQCPEHSQLTTLGVDGKEFPEVHLGQWYFIAGAAPTKEELATFDPVDNIVFNMAA

GSAPMQLHLRATIRMKDGLCVPRKWIYHLTEGSTDLRTEGRPDMKTELFSSSCPGGIMLNETGQGYQRFLLYN

RSPHPPEKCVEEFKSLTSCLDSKAFLLTPRNQEACELSNNVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.