

32-7519: Recombinant Human CD3 e/CD3E (C-6His)(Discontinued)

Gene : CD3E
Gene ID : 916
Uniprot ID : P07766

Description

Source: Human Cells.

MW :12.79kD.

Recombinant Human CD3 epsilon is produced by our Mammalian expression system and the target gene encoding Asp23-Asp126 is expressed with a 6His tag at the C-terminus. T-Cell Surface Glycoprotein CD3 e Chain (CD3e) is a single-pass type I membrane protein. CD3e contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3e is a polypeptide encoded by the CD3E gene on chromosome 11 in humans. The T cell receptor-CD3 complex (TCR/CD3 complex) is involved in T-cell development and several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs). TCR/CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : DGNEEMGGITQTPYKVSISGTTVILTCPQYPGSEILWQHNDKNIGGEDDKNIGSDEDHLSLKEFSELEQSGYYVCYPRGSKPEDANFYLYLRVCENCMEMDVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.