

32-7526: Recombinant Human Complement Factor H-Related 1/CFHR1 (C-6His)

Gene : CFHR1
Gene ID : 3078
Uniprot ID : Q03591

Description

Source: Human Cells.
MW :36.78kD.

Recombinant Human CFHR1 is produced by our Mammalian expression system and the target gene encoding Glu19-Arg330 is expressed with a 6His tag at the C-terminus. Complement Factor H-Related 1 (CFHR1) is a 43 kDa secreted member of the factor H family of glycoproteins. The human Complement Factor H protein family consists of the complement and immune regulators factor H, the factor H-like protein 1 (FHL-1) and five factor H-related proteins (CFHR-1 to -5). Members of the H-related protein family are exclusively composed of individually folded protein domains, termed short consensus repeats (SCRs) or complement control modules. FHR1 is produced by hepatocytes and circulates as two differentially glycosylated isoforms (37 kDa and 43 kDa). Mature human FHR1 is 312 amino acids in length. It contains five, approximately 60 aa SCRs that basically constitute the entire molecule. FHR1 may play a role in complement regulation, lipid metabolism and lipoprotein complexes that bind PMNs to LPS.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : EATFCDFPKINHGILYDEEKYKPFVQVPTGEVFFYSCEYNFVSPSKSFWTRITCTEEGWSPTPKCLRLCFFPFVEN
GHSESSGQTHLEGDTVQIICNTGYRLQNNENNISCVERGWSTPPKCRSTDTSCVNPPTVQNAHILSRQMSKYP
SGERVRYECSRSPYEMFGDEEVMCLNGNWTEPPQCKDSTGKCGPPPIDNGDITSFPLSVYAPASSVEYQCQNL
YQLEGNKRITCRNGQWSEPPKCLHPCVISREIMENYNIALRWTAKQKLYLRTGESAEFVCKRGYRLSSRSHTLRT
TCWDGKLEYPTCAKRDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.