

## 32-7532: Recombinant Human Carboxypeptidase A1/CPA1 (C-6His)(Discontinued)

**Gene :** CPA1  
**Gene ID :** 1357  
**Uniprot ID :** P15085

### Description

Source: Human Cells.  
MW :46.57kD.

Recombinant Human Carboxypeptidase A1 is produced by our Mammalian expression system and the target gene encoding Lys17-Tyr419 is expressed with a 6His tag at the C-terminus. Carboxypeptidase A1 (CPA1) is secreted as a pancreatic peptidase that comes from the precursor form of inactive procarboxypeptidase. CPA1 comprises a signal peptide, a pro region and a mature chain, and can be activated after cleavage of the pro peptide. It has a free C-terminal carboxyl group, with the preference of residues with aromatic or branched aliphatic side chains. CPA1 cleaves the C-terminal amide or ester bond of peptides and involves in zymogen inhibition. Three different forms of human pancreatic procarboxypeptidase A have been isolated. In contrast to procarboxypeptidase B which was always secreted by the pancreas as a monomer, procarboxypeptidase A occurs as a monomer and/or associated to one or two functionally different proteins, such as zymogen E.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Supplied as a 0.2 µm filtered solution of 20mM TrisHCl, 150mm NaCl, pH 7.5.  
**Storage condition :** Store at -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.  
**Amino Acid :** KEDFVGHQVLRISVADEAQVQKVELEDLEHLQLDFWRGPAHPGSPIDVRVPFPSIQAVKIFLESHGISYETMIE  
DVQSLLEDDEEQEMFAFRSRARSTDTFNATYHTLEEIYDFLDLLVAENPHLVSKIQIGNTYEGRPYVLFKSTGGS  
KRPAIWIDTGIHSREWVTQASGVVFAKKITQDYGQDAAFTALDLDIFLEIVTNPDGFAFTHSTNRMWRKTRS  
HTAGSLCIGVDPNRRWDAGFGLSGASSNPCSETYRGKFANSEVEVKSIVDFVKDHGNIKAFISHSYSQLLMYPY  
GYKTEPVPDQDELQLSKAAVTALASLYGTFKFNYSIIKAIYQASGSTIDWTYSQGIKYSFTFELRDTGRYGFLLP  
ASQIIPTAKETWLALLTIMEHTLNHPYVDHHHHHH

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.