

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7541: Recombinant Human Glioma Pathogenesis-Related Protein 1/GLIPR1/RTVP1 (C-6His)

Gene : GLIPR1
Gene ID : 11010
Uniprot ID : P48060

Description

Source: Human Cells. MW:25.16kD.

Recombinant Human GLIPR1 is produced by our Mammalian expression system and the target gene encoding Asn22-Arg232 is expressed with a 6His tag at the C-terminus. Glioma Pathogenesis-Related Protein 1 (GLIPR1) is a member of the CRISP family. It is a single-pass membrane protein with 266 amino acids. GLIPR1 is highly expressed in the lung and kidney and lowly expressed in the heart and liver. It is also highly expressed in cell lines that are derived from nervous system tumors arising from glia; conversely, it is lowly expressed in non-glial-derived nervous system tumor cell lines. GLIPR1 is only expressed in brain tumor glioblastoma multiforme/astrocytoma and not expressed in other nervous system tumors nor normal adult or fetal tissues.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: ANILPDIENEDFIKDCVRIHNKFRSEVKPTASDMLYMTWDPALAQIAKAWASNCQFSHNTRLKPPHKLHPNFTSL

GENIWTGSVPIFSVSSAITNWYDEIQDYDFKTRICKKVCGHYTQVVWADSYKVGCAVQFCPKVSGFDALSNGA HFICNYGPGGNYPTWPYKRGATCSACPNNDKCLDNLCVNRQRDQVKRYYSVVYPGWPIYPRNRVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.