

## 32-7551: Recombinant Human Nectin-3/PVRL3/CD113 (C-6His)

**Gene :** NECTIN3

**Gene ID :** 25945

**Uniprot ID :** Q9NQS3

### Description

Source: Human Cells.

MW :34.96kD.

Recombinant Human Nectin-3 is produced by our Mammalian expression system and the target gene encoding Gly58-Cys366 is expressed with a 6His tag at the C-terminus. Nectin-3 is a type I transmembrane glycoprotein that belongs to the nectin family. Its precursor is 549 amino acids in length and contains an extended signal sequence of 57 amino acids, an extracellular domain (ECD) of 347 amino acids, a transmembrane segment of 21 amino acids, and a cytoplasmic region of 124 amino acids. It is predominantly expressed in testis and placenta as well as in various cell lines, including epithelial cell lines. Nectin-3 plays a role in cell-cell adhesion through heterophilic trans-interactions with nectin-like proteins or nectins, such as trans-interaction with PVRL2/Nectin-2 at Sertoli-spermatid junctions. Nectin-3 is also involved in the formation of cell-cell junctions, including adherens junctions and synapses.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** GPIIVEPHVTAVWGKNVSLKCLIEVNETITQISWEKIHGKSSQTVAVHHPQYGFVSVQGEYQGRVLFKNYSLN DAT  
ITLHNIGFSDSGKYICKAVTFPLGNAQSSTTVTLVEPTVSLIKGPD SLIDGGNETVAAICIAATGKPV AHIDWEGD  
LGEMESTTTSFPNETATIISQYKLPTRFARGRRITCVVKHQALEKDIRYSFILD IQYAPEVSVTG YDGNW FVGRK  
GVNLKCNADANPPPFSVWSRLDGQWPDG LLASDNTLHFVHPLTFNYSGVYICKVTNSL GQRSDQKV IYISAY  
NSVASLNCVDHHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.