

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7612: Recombinant Mouse Interleukin-11/IL-11 (C-6His)(Discontinued)

Gene ID: 16156 **Uniprot ID:** P47873

Description

Source: Human Cells. MW :20.2kD.

Recombinant Mouse Interleukin-11 is produced by our Mammalian expression system and the target gene encoding Pro22-Leu199 is expressed with a 6His tag at the C-terminus. Interleukin-11(IL-11) is a secreted protein and belongs to the IL-6 superfamily. IL-11 has been demonstrated to improve platelet recovery after chemotherapy-induced thrombocytopenia, induceacute phase proteins, modulate antigen-antibody responses, participate in the regulation of bone cellproliferation and differentiation and could be use as a therapeutic for osteoporosis. IL-11 stimulates the growth of certain lymphocytes and, in the murine model, stimulates an increase in the cortical thickness and strength of long bones. In addition to having lymphopoietic/hematopoietic and osteotrophic properties, it has functions in many other tissues, including the brain, gut, testis and bone.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: PGPPAGSPRVSSDPRADLDSAVLLTRSLLADTRQLAAQMRDKFPADGDHSLDSLPTLAMSAGTLGSLQLPGVL

TRLRVDLMSYLRHVQWLRRAGGPSLKTLEPELGALQARLERLLRRLQLLMSRLALPQAAPDQPVIPLGPPASAW

GSIRAAHAILGGLHLTLDWAVRGLLLLKTRLVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.